

NOTE ON q -DEDEKIND TYPE SUMS RELATED TO q -EULER POLYNOMIALS

TAEKYUN KIM

ABSTRACT. Recently, q -Dedekind type sums related to q -zeta function and basic L -series are studied by Simsek in [Y. Simsek, q -Dedekind type sums related to q -zeta function and basic L -series, J. Math. Anal. Appl. 318(2006) 333-351] and Dedekind type sums related to Euler numbers and polynomials are introduced in previous paper [T. Kim, Note on Dedekind type DC sums, Adv. Stud. Contemp. Math. 18(2009), 249-260]. It is the purpose of this paper to construct a p -adic continuous function for an odd prime to contain a p -adic q -analogue of higher order Dedekind type sums related to q -Euler polynomials and numbers by using an invariant p -adic q -integrals.

§1. Introduction/ Preliminaries

Let p be a fixed odd prime number. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field and the completion of algebraic closure of $\mathbb{Q}_p$. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = \frac{1}{p}$. When one talks of q -extension, q is variously considered as and indeterminate, a complex number $q \in \mathbb{C}$ or p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one normally assumes $|q| < 1$. If $q \in \mathbb{C}_p$, one normally assumes $|1 - q|_p < 1$. Recently, we proposed a definition of a q -extension of p -adic Haar measure as follows: For any positive integer N , we set

$$\mu_q(a + p^N \mathbb{Z}_p) = \frac{(1+q)(-q)^a}{1+q^{p^N}}, \quad (\text{see [1-12]}).$$

for $0 \leq a \leq p^N - 1$ and this can be extended to a measure on $\mathbb{Z}_p$. This measure yields an invariant p -adic q -integral for each non-negative integer m and the m -th Carlitz's type q -Euler numbers $\varepsilon_{n,q}$ can be represented by this p -adic q -integral as follows:

$$\varepsilon_{n,q} = \int_{\mathbb{Z}_p} \left(\frac{1-q^a}{1-q} \right)^m d\mu_q(a) = \lim_{N \rightarrow \infty} \frac{1+q}{1+q^{p^N}} \sum_{x=0}^{p^N-1} \left(\frac{1-q^a}{1-q} \right)^m (-q)^a,$$

Key words and phrases. : DC sums, Dedekind sums, Euler numbers, p -adic q -Dedekind sums.

2000 AMS Subject Classification: 11B68, 11S80

The present Research has been conducted by the research Grant of Kwangwoon University in 2008

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -T $\mathcal{E}\mathcal{X}$

which has a sense as we see readily that the limit is convergent (see[8,9]). The modified q -Euler numbers are also defined as

$$E_{m,q} = \int_{\mathbb{Z}_p} \left(\frac{1-q^a}{1-q} \right)^m q^{-a} d\mu_q(a), \text{ (see [6])}.$$

Note that $\lim_{q \rightarrow 1} E_{n,q} = E_n$, where E_n are the n -th Euler numbers. Now, we also consider the q -Euler polynomials as follows:

$$E_{m,q}(x) = \int_{\mathbb{Z}_p} q^{-t} \left(\frac{1-q^{x+t}}{1-q} \right)^m d\mu_q(t), \text{ for } x \in \mathbb{Z}_p, m \in \mathbb{N}.$$

These numbers $E_{m,q}(x)$ can be represented by

$$E_{m,q}(x) = \sum_{l=0}^m \binom{m}{l} q^{xl} E_{l,q} \left(\frac{1-q^x}{1-q} \right)^{m-l}.$$

For any positive integer h, k and m , Dedekind type DC sums are defined as

$$S_m(h, k) = \sum_{M=1}^{k-1} (-1)^{M-1} \frac{M}{k} \bar{E}_m\left(\frac{hM}{k}\right), \text{ (see [7]),}$$

where $\bar{E}_m(x)$ are the m -th periodic Euler function.

By using an invariant p -adic q -integral on $\mathbb{Z}_p$, we construct a p -adic continuous function for an odd prime to contain a p -adic q -analogue of higher order Dedekind type DC sums $k^m S_{m+1}(h, k)$ in this paper. It is the purpose of this paper to give a q -analogue of p -adic Dedekind type DC sums by using invariant p -adic q -integral on $\mathbb{Z}_p$ approach the p -adic analogue of higher order Dedekind sums at $q = 1$ as follows:

Theorem. *Let h, k be positive integer with $(h, k) = 1$, $p \nmid k$. For $s \in \mathbb{Z}_p$, let us define p -adic Dedekind type DC sums as follows.*

$$S_{p,q}(s : h, k : q^k) = \sum_{M=1}^{k-1} \left(\frac{1-q^M}{1-q} \right) (-1)^{M-1} T_q(s, hM, k : q^k).$$

Then there exists a continuous function $S_{p,q}(s : h, k : q^k)$ on $\mathbb{Z}_p$ which satisfies

$$\begin{aligned} S_{p,q}(m : h, k : q^k) &= \left(\frac{1-q^k}{1-q} \right)^{m+1} S_{m,q}(h, k : q^k) \\ &\quad - \left(\frac{1-q^k}{1-q} \right)^{m+1} \left(\frac{1-q^{kp}}{1-q^k} \right)^m S_{m,q}((p^{-1}h)_k, k : q^{pk}), \end{aligned}$$

where $m+1 \equiv 0 \pmod{p-1}$, and $(p^{-1}a)_N$ denotes the integer x with $0 \leq x < N$, $px \equiv a \pmod{N}$.

§2. Proof of theorem

The q -Euler numbers $E_{m,q}$ can be written as

$$E_{0,q} = \frac{1+q}{2}, \quad (qE+1)^n + E_{n,q} = 0 \text{ if } n > 0,$$

which is

$$E_{n,q} = (1+q) \left(\frac{1}{1-q} \right)^n \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+q^l}, \quad (\text{see [6]}),$$

where we use the technique method notation by replacing E^n by $E_{n,q}$, symbolically. Let w denote Teichüller character $(\text{mod } p)$. For $x \in \mathbb{Z}_p^\times = \mathbb{Z}_p \setminus p\mathbb{Z}_p$, we set

$$\langle x \rangle_q = \langle x : q \rangle = w^{-1}(x) \left(\frac{1-q^x}{1-q} \right).$$

Let a and N be positive integers with $(p, a) = 1$ and $p|N$. Define

$$T_q(s, a, N : q^N) = w^{-1}(a) \langle a \rangle_q^s \sum_{j=0}^{\infty} \binom{s}{j} q^{aj} \left(\frac{1-q^N}{1-q^a} \right)^j E_{j,q^N}, \quad \text{for } s \in \mathbb{Z}_p.$$

In particular, if $m+1 \equiv 0 \pmod{p-1}$, then

$$\begin{aligned} T_q(m, a, N : q^N) &= \left(\frac{1-q^a}{1-q} \right)^m \sum_{j=0}^m \binom{m}{j} q^{aj} \left(\frac{1-q^N}{1-q^a} \right)^j E_{j,q^N} \\ &= \left(\frac{1-q^N}{1-q} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1-q^{Nx+a}}{1-q^N} \right)^m q^{-Nx} d\mu_{q^N}(x). \end{aligned}$$

Therefore, $T_q(m, a, N : q^N)$ is a continuous p -adic extension of $\left(\frac{1-q^N}{1-q} \right)^m E_{m,q^N}(\frac{a}{N})$.

Let $[\cdot]$ be Gauss' symbol and let $\{x\} = x - [x]$. Then we now consider a q -analogue of higher order Dedekind type DC sums $S_{m,q}(h, k : q^l)$ as follows:

$$S_{m,q}(h, k : q^l) = \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1-q^M}{1-q^k} \right) \int_{\mathbb{Z}_p} q^{-lx} \left(\frac{1-q^{l(x+\{\frac{hM}{k}\})}}{1-q^l} \right)^m d\mu_{q^l}(x).$$

If $m+1 \equiv 0 \pmod{p-1}$, then we have

$$\begin{aligned} (1) \quad & \left(\frac{1-q^k}{1-q} \right)^{m+1} \sum_{M=1}^{k-1} \left(\frac{1-q^M}{1-q^k} \right) (-1)^{M-1} \int_{\mathbb{Z}_p} \left(\frac{1-q^{k(x+\{\frac{hM}{k}\})}}{1-q^k} \right)^m q^{-kx} d\mu_{q^{-k}}(x) \\ &= \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1-q^M}{1-q} \right) \left(\frac{1-q^k}{1-q} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1-q^{k(x+\{\frac{hM}{k}\})}}{1-q^k} \right)^m q^{-kx} d\mu_{q^k}(x), \end{aligned}$$

where $p|k$, $(hM, p) = 1$ for each M . From (1), we note that

$$\begin{aligned}
& \left(\frac{1-q^k}{1-q} \right)^{m+1} S_{m,q}(h, k : q^k) \\
(2) \quad &= \sum_{M=1}^{k-1} \left(\frac{1-q^M}{1-q} \right) \left(\frac{1-q^k}{1-q} \right)^m (-1)^{M-1} \int_{\mathbb{Z}_p} q^{-kx} \left(\frac{1-q^{k(x+\frac{hM}{k})}}{1-q^k} \right)^m d\mu_{q^k}(x) \\
&= \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1-q^M}{1-q} \right) T_q(m, (hM)_k : q^k),
\end{aligned}$$

where $(y)_k$ denotes the integers x such that $0 \leq x < n$ and $x \equiv \alpha \pmod{k}$.

It is easy to check that

$$\begin{aligned}
& \int_{\mathbb{Z}_p} q^{-t} \left(\frac{1-q^{x+t}}{1-q} \right)^k d\mu_q(t) \\
(3) \quad &= \left(\frac{1-q^m}{1-q} \right)^k \frac{1+q}{1+q^m} \sum_{i=0}^{m-1} (-1)^i \int_{\mathbb{Z}_p} \left(\frac{1-q^{m(t+\frac{x+i}{m})}}{1-q^m} \right)^k q^{-mt} d\mu_{q^m}(t).
\end{aligned}$$

By (2) and (3), we see that

$$\begin{aligned}
& \left(\frac{1-q^N}{1-q} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1-q^{N(x+\frac{a}{N})}}{1-q^N} \right)^m q^{-Nx} d\mu_{q^N}(x) \\
(4) \quad &= \frac{1+q^N}{1+q^{pN}} \sum_{i=0}^{p-1} \left(\frac{1-q^{Np}}{1-q} \right)^m (-1)^i \int_{\mathbb{Z}_p} \left(\frac{1-q^{pN(x+\frac{a+iN}{pN})}}{1-q^{pN}} \right)^m q^{-pNx} d\mu_{q^{pN}}(x).
\end{aligned}$$

From (2), (3) and (4) we note that the p -adic integration is given by

$$T_q(s, a, N : q^N) = \frac{1+q^N}{1+q^{pN}} \sum_{\substack{i=0 \\ a+iN \not\equiv 0 \pmod{p}}}^{p-1} (-1)^i T_q(s, (a+iN)_{pN}, p^N : q^{pN})$$

such that

$$\begin{aligned}
& T_q(m, a, N : q^N) \\
&= \left(\frac{1-q^N}{1-q} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1-q^{N(x+\frac{a}{N})}}{1-q^N} \right)^m q^{-Nx} d\mu_{q^N}(x) \\
&- \left(\frac{1-q^{pN}}{1-q} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1-q^{pN(x+\frac{(p-1)a}{N})}}{1-q^{pN}} \right)^m q^{-pNx} d\mu_{q^{pN}}(x),
\end{aligned}$$

where $(p^{-1}a)_N$ denotes the integer x with $0 \leq x < N$, $px \equiv a \pmod{N}$ and m is integer with $m+1 \equiv 0 \pmod{p-1}$.

Hence, we have

$$(5) \quad \sum_{M=1}^{k-1} \left(\frac{1-q^M}{1-q} \right) (-1)^{M-1} T_q(m, hM, k : q^k) = \left(\frac{1-q^k}{1-q} \right)^{m+1} S_{m,q}(h, k : q^k) \\ - \left(\frac{1-q^k}{1-q} \right)^{m+1} \left(\frac{1-q^{kp}}{1-q^k} \right)^m S_{m,q}((p^{-1}h)_k, k : q^{pk}),$$

where $p \nmid k$ and $p \nmid hM$ for each M .

For $s \in \mathbb{Z}_p$, let us define p -adic Dedekind type DC sums as follows:

$$S_{p,q}(s : h, k : q^k) = \sum_{M=1}^{k-1} \left(\frac{1-q^M}{1-q} \right) (-1)^{M-1} T_q(s, hM, k : q^k).$$

Then there exists a continuous function $S_{p,q}(s : h, k : q^k)$ on $\mathbb{Z}_p$ which satisfies

$$S_{p,q}(m : h, k : q^k) = \left(\frac{1-q^k}{1-q} \right)^{m+1} S_{m,q}(h, k : q^k) \\ - \left(\frac{1-q^k}{1-q} \right)^{m+1} \left(\frac{1-q^{kp}}{1-q^k} \right)^m S_{m,q}((p^{-1}h)_k, k : q^{pk}), \text{ where } m+1 \equiv 0 \pmod{p-1}.$$

Remark. Note that

$$S_m(h, k : q^l) = \sum_{M=1}^{k-1} \left(\frac{1-q^M}{1-q^k} \right) (-1)^{M-1} E_{m,q^l}(\{ \frac{hM}{k} \}).$$

It is easy to see that $S_{p,1}(s : h, k : 1)$ is the p -adic analogue of higher order Dedekind type DC sums $k^m S_{m+1}(h, k)$.

REFERENCES

- [1] T. M. Apostol, *Generalized Dedekind sums and transformation formulae of certain Lambert series*, Duke Math. J. **17** (1950), 147-157.
- [2] B. C. Berndt, *Dedekind sums and paper of G. H. Hardy*, J. London Math. Soc. **13** (1976), 129-136.
- [3] B. C. Berndt, *Generalized Dedekind eta functions and generalized Dedekind sums*, Trans. Amer. Math. Soc. **178** (1973), 495-508.
- [4] M. Can, M. Cenkci, V. Kurt, Y. Simsek,, *Twisted Dedekind type sums associated with Barnes' type multiple Frobenius-Euler l -functions*, Adv. Stud. Contemp. Math. **18** (2009), 135-160.
- [5] M. Cenkci, M. Can and V. Kurt, *p -adic interpolation functions and Kummer-type congruences for q -twisted Euler numbers*, Adv. Stud. Contemp. Math. **9** (2004), 203-216.

- [6] T. Kim, *The modified q -Euler numbers and polynomials*, Adv. Stud. Contemp. Math. **16** (2008), 161-170.
- [7] T. Kim, *Note on Dedekind type DC sums*, Adv. Stud. Contemp. Math. **18** (2009), 249-260.
- [8] T. Kim, *q -Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
- [9] T. Kim, *On p -adic interpolating function for q -Euler numbers and its derivatives*, J. Math. Anal. Appl. **339** (2008), 598-608.
- [10] T. Kim, *q -Euler numbers and polynomials associated with p -adic q -integrals*, J. Nonlinear Math. Phys. **14** (2007), 15-27.
- [11] Y. Simsek, *q -Dedekind type sums related to q -zeta function and basic L -series*, J. Math. Anal. Appl. **318** (2006), 333-351.
- [12] Y. Simsek, *Generalized Dedekind sums associated with the Abel sums and the eisenstein and Lambert series*, Adv. Stud. Contemp. Math. **9** (2004), 195-202.

Taekyun Kim

Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, S. Korea e-mail: tkkim@kw.ac.kr